

PEEL FROM CORNER OF BOOKLET

15M-27-29K

Distributed By:
VOLUNTARY PURCHASING GROUPS, INC.
 230 FM 87
 BONHAM, TEXAS 75418
 Visit Us At: www.fertilome.com
 EPA Reg. No. 2217-597-7401
 EPA Est. No. 7401-Tx-001




FIRST AID


If In Eyes:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a Poison Control Center or doctor for treatment advice.
If On Skin Or Clothing:	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a Poison Control Center or doctor for treatment advice.
If Swallowed:	<ul style="list-style-type: none"> • Call a Poison Control Center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by a Poison Control Center or doctor. • Do not give anything by mouth to an unconscious person.
If Inhaled:	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. • Call a Poison Control Center or doctor for further treatment advice.


Have the product container or label with you when calling a Poison Control Center or doctor or going for treatment. For additional information in case of emergency, call toll free 1-800-858-7378 or 1-800-222-1222.

Note To Physician: Probable mucosal damage may contraindicate the use of gastric lavage.




ferti·lome®







DANDELION



CHICKWEED



CLOVER



Weed-Out®

Lawn Weed Killer

- Contains TRIMEC® Herbicide.
- Kills Troublesome Weeds: Dandelion, Chickweed, Clover And Many Others.

ACTIVE INGREDIENTS:	
DIMETHYLAMINE SALT OF (+)-(R)-2-(2-METHYL-4-CHLOROPHENOXY) PROPIONIC ACID.....	5.45%
DIMETHYLAMINE SALT OF 2,4-DICHLOROPHENOXYACETIC ACID.....	5.88%
DIMETHYLAMINE SALT OF DICAMBA (3,6-DICHLORO-O-ANISIC ACID).....	1.21%
INERT INGREDIENTS:	87.46%
TOTAL:	100.00%

THIS PRODUCT CONTAINS:
 0.39 lb. (+)-(R)-2-(2-methyl-4-chlorophenoxy) propionic acid equivalent per gallon or 4.50%.
 0.42 lb. 2,4-dichlorophenoxyacetic acid equivalent per gallon or 4.88%.
 0.09 lb. 3,6-dichloro-o-anisic acid equivalent per gallon or 1.00%.
 Isomer Specific by AOAC Methods.
 Contains the single isomer form of Mecoprop-p.
 TRIMEC® is a registered trademark of PBI/GORDON CORPORATION.

KEEP OUT OF REACH
 OF CHILDREN
ANGER
 See Back Panel And Attached Booklet
 For Additional Precautionary
 Statements, First Aid And
 Directions for Use.

NET CONTENTS ONE PINT (16 FL. OZS.)



READ THE ENTIRE LABEL FIRST.

OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

DANGER: Corrosive. Causes eye damage. Do not get in eyes or on skin or clothing. Wear goggles when applying this product. Harmful if swallowed or inhaled. Avoid exposure to spray mist. When using this product, wear long-sleeved shirt, long pants, socks, shoes, rubber gloves and eye protection. It is recommended that safety glasses include front, brow and temple protection.

After using this product, rinse gloves before removing, remove clothing and laundry separately before reuse, and promptly and thoroughly wash hands and exposed skin with soap and water. Remove saturated clothing as soon as possible and shower.

ENVIRONMENTAL HAZARDS

This product is toxic to aquatic invertebrates. Drift or runoff may adversely affect aquatic invertebrates and nontarget plants. Do not apply directly to water. Do not contaminate water when disposing of equipment washwater. Do not apply when weather conditions favor drift from target area.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Do not allow people (other than applicator) or pets on treatment area during application. Do not enter treatment area until spray has dried or dust has settled.

WHEN TO USE

Spray at any time when Weeds are growing. For best results, apply in Spring when Weeds are small and growing rapidly. Apply in Fall to remove late germinating Weeds.

HOW TO USE

The maximum application rate to turf is 0.6 pounds of 2,4-D acid equivalent per acre per application per site. The maximum number of broadcast applications per treatment site is 2 per year.

Hose-End Sprayer: Use 1 gallon of water for every 250 square feet to be sprayed. For concentrate sprayers, add only the required amount of **ferti-lome® WEED-OUT** listed in the rate chart depending on type of Grass and number of square feet to be treated. Do not add water to concentrate sprayers.

For dilution sprayers, add the required amount of **ferti-lome® WEED-OUT** and fill sprayer with water to proper level. Spray uniformly.

Pressure-Sprayer: Use ½ gallon of water for every 250 square feet to be sprayed. Add the required amount of **ferti-lome® WEED-OUT** listed in the rate chart for the type of Grass to be treated and fill sprayer with water to the proper level. Adjust spray nozzle to deliver coarse spray. Spray uniformly.

Spot-Treatment: If hardy Weeds persist in Sensitive Grasses such as St. Augustine, Centipedegrass, Bermudagrass, Buffalograss or Bentgrass, spot treat individual Weeds using 1 teaspoon of **ferti-lome® WEED-OUT** in 1 quart of water (equivalent to 4 teaspoons per gallon of water.) Minimize contact of Grass.

RATES OF FERTI-LOME® WEED-OUT FOR VARIOUS LAWN GRASSES					
Gallons Of Water Spray		Area to Treat (Sq. Ft.)	St. Augustinegrass, Centipedegrass (fl. oz.)	Bermudagrass, Buffalograss Bentgrass	Bluegrass Zoysiagrass, Fescue (fl. oz)
Hose End Sprayer	Pressure Sprayer				
1	½	250	½	2/3	2/3 to 1
4	2	1,000	2	2 2/3	2 2/3 to 4
8	4	2,000	4	5 ½	5 ½ to 8
20	10	5,000	10	13 1/3	13 1/3 to 20
32	16	8,000	16	21 1/3	21 1/3 to 32

EQUAL MEASURES

1 fluid ounce (fl. oz.) = 2 tablespoons (Tbs.) or 6 teaspoons (tsp.)
 2 fluid ounces = 4 tablespoons 4 fluid ounces = ½ cup 8 fluid ounces = 1 cup
 16 fluid ounces = 1 pint (pt.) 32 fluid ounces = 1 quart (qt.)

HOW TO USE IN THE THE 20 GALLON HOSE-END SPRAYER

1. Connect sprayer nozzle to garden hose; remove safety cap from container and insert sprayer unit, securing tightly.
2. Turn on water; place finger over hole on right side of nozzle and product will automatically be dispersed.
3. Spray evenly and uniformly over area requiring application. (Check application guide to determine the correct area to be covered per ounce.)

***APPLICATION GUIDE**

Fluid Ounces	Recommended Coverage Area Per Ounce		
	Sensitive Grasses Such As Centipedegrass And St. Augustinegrass (sq.ft.)	Bermudagrass, Bentgrass, And Buffalograss (sq. ft.)	Bluegrass, Fescue, And Zoysiagrass (sq. ft.)
1	500	375	250
2	1,000	750	500
3	1,500	1,125	750
4	2,000	1,500	1,000
5	2,500	1,875	1,250
6	3,000	2,250	1,500
7	3,500	2,525	1,750
8	4,000	3,000	2,000
9	4,500	3,375	2,250
10	5,000	3,750	2,500
11	5,500	4,125	2,750
12	6,000	4,500	3,000
13	6,500	4,875	3,250
14	7,000	5,250	3,500
15	7,500	5,625	3,750
16	8,000	6,000	4,000

*Good for this precalibrated sprayer only.

Use Precautions: Do not apply this product through any type of irrigation system. Use only lawn sprayer that delivers coarse spray to reduce drift. Avoid spray drift to Vegetables, Flowers, Ornamental Plants. Do not spray roots of Trees and Ornamentals. Do not exceed specified dosages for any area; be particularly careful within dripline of Trees and other Ornamental species. Do not spray Carpetgrass, Dichondra, or Lawn where desirable Clovers are present. Do not apply this product to "Floritam" St. Augustine Grass in Florida. Do not apply to newly seeded Lawn Grasses until well established. Do not spray when air temperatures exceed 85°F. Seed can be sown 3 to 4 weeks after application. After using this product, clean sprayer with soap or detergent and water, and rinse thoroughly before applying other pesticides. Failure to follow above precautions may result in injury.

GENERAL INFORMATION

FERTI-LOME® WEED-OUT controls troublesome common Lawn Weeds, including Poison Ivy, in both Northern and Southern Grasses.

- BROADLEAF WEEDS -

Annual Yellow	Buckhorn	Clover, Sweet	Hairy Fleabane
Sweetclover	Buckhorn Plantain	Clover, White	Hawkweed
Aster	Bulbous Buttercup	Cockle	Healall
Austrian Fieldcress	Bull Thistle	Cocklebur	Heartleaf Drymary
Bedstraw	Bullnettle	Common Mullein	Heath Aster
Beggarticks	Burclover	Creeping Jenny	Hedge Bindweed
Betony, Florida	Burdock	Cudweed	Hedge Mustard
Bindweed, Field	Burning Nettle	Curly Dock	Henbit
Bird Vetch	Burweed	Daisy, English	Hoary Cress
Bitter Wintercress	Buttercup	Daisy Fleabane	Hoary Plantain
Bittercress, Hairy	Buttonweed	Daisy, Oxeye	Hoary Vervain
Bitterweed	Canada Thistle	Dandelion	Horsenettle
Black-Eyed Susan	Carolina Geranium	Dichondra	Jimsonweed
Black Medic	Carpetweed	Dogfennel	Knawel
Black Mustard	Catchweed Bedstraw	Dollarweed	Knotweed
Blackseed Plantain	Catnip	False Dandelion	Kochia
Blessed Thistle	Catsear	Falseflax	Lambsquarters
Bloodflower Milkweed	Chickweed, Common	False Sunflower	Lespedeza
Blue Lettuce	Chickweed, Mouseear	Fiddleneck	Mallow
Blue Vervain	Chicory	Florida Pusley	Matchweed
Bracted Plantain	Cinquefoil	Frenchweed	Mexicanweed
Bristly Oxtongue	Clover, Crimson	Galinsoga	Milk Vetch
Broadleaf Dock	Clover, Hop	Goldenrod	Morningglory
Broadleaf Plantain	Clover, Red	Ground Ivy	Mouseear Hawkweed
Broomweed	Clover, Strawberry	Gumweed	Mugwort

- BROADLEAF WEEDS - (Continued)

Musk Thistle	Prostrate Pigweed	Spiny Cocklebur	Wild Buckwheat
Mustard	Prostrate Spurge	Spiny Sowthistle	Wild Carrot
Narrowleaf Plantain	Prostrate Vervain	Spotted Catsear	Wild Four-O'Clock
Narrowleaf Vetch	Puncturevine	Spotted Spurge	Wild Garlic
Nettle	Purslane, Common	Spurweed	Wild Geranium
Orange Hawkweed	Ragweed	Stinging Nettle	Wild Lettuce
Oriental Cocklebur	Red Sorrel	Strawberry, India Mock	Wild Marigold
Oxalis	Redroot Pigweed	Tall Nettle	Wild Mustard
Parsley-Piert	Redstem Filaree	Tall Vervain	Wild Onion
Parsnip	Rough Cinquefoil	Tansy Ragwort	Wild Parsnip
Pearlwort	Rough Fleabane	Tansy Mustard	Wild Radish
Pennycress	Russian Pigweed	Tanweed	Wild Rape
Pennywort	Russian Thistle	Thistle	Wild Strawberry
Pepperweed	Scarlet Pimpernel	Trailing Crownvetch	Wild Sweet Potato
Pigweed	Scotch Thistle	Tumble Mustard	Wild Vetch
Pineywoods Bedstraw	Sheep Sorrel	Tumble Pigweed	Woodsorrel
Plains Coreopsis	Sherherdspurse	Velvetleaf	Woolly Croton
Plantain	Slender Plantain	Venice Mallow	Woolly Morningglory
Poison Ivy	Smallflower Galinsoga	Virginia Buttonweed	Woolly Plantain
Poison Oak	Smooth Dock	Virginia Creeper	Wormseed
Pokeweed	Smooth Pigweed	Virginia Pepperweed	Yarrow
Poorjoe	Sorrel	Wavyleaf Bullthistle	Yellow Rocket
Prairie Sunflower	Sowthistle	Western Clematis	Yellowflower
Prickly Lettuce	Spanishneedles	Western Salsify	Pepperweed
Prickly Sida	Speedwell	White Mustard	and other Broadleaf
Prostrate Knotweed	Spiny Amaranth	Wild Aster	Weeds

STORAGE AND DISPOSAL

STORAGE: Keep from freezing. Store in original container and place in locked storage area when not using.

CONTAINER DISPOSAL: If Empty: Do not reuse this container. Place in trash or offer for recycling if available. **If Partially Filled:** Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

LIMITED WARRANTY AND DISCLAIMER

The Manufacturer warrants only that the chemical composition of this product conforms to the ingredient statement given on the label and that the product is reasonably suited for the labeled use when applied according to the Directions for Use.

THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY OTHER EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE EXPRESSLY DISCLAIMED. This Limited Warranty does not extend to the use of the product inconsistent with label instructions, warnings or cautions, or to use of the product under abnormal conditions such as drought, excessive rainfall, tornadoes, hurricanes, etc. These factors are beyond the control of the Manufacturer or the Seller. Any damages arising from a breach of the Manufacturer's Warranty shall be limited to direct damages, and shall not include indirect or consequential damages such as loss or profits or values, except as otherwise provided by law.

The terms of this Limited Warranty and Disclaimer cannot be varied by any written or verbal statements or agreements. No employee or agent of the Seller is authorized to vary or exceed the terms of this Limited Warranty and Disclaimer in any manner.

Distributed By:
VOLUNTARY PURCHASING GROUPS, INC.

230 FM 87
BONHAM, TEXAS 75418
EPA Reg. No. 2217-597-7401
EPA Est. No. 7401-TX-001
Visit Us At: www.fertilome.com

