

33408

WEED & FEED

# Hi-Yield® Weed & Feed



**Controls Over 200 Listed Broadleaf Weeds, Roots And All, Including Dandelion, Chickweed, Plantain, Knotweed, Henbit And Spurge.**

- **CONTROLS:** Dandelion, Chickweed, Knotweed, Plantain, Henbit, Spurge And Many Other Broadleaf Weeds In St. Augustinegrass, Centipedegrass, Bahiagrass, Bermudagrass, Zoysiagrass, Bluegrass, Perennial Ryegrass, Fescue *spp.* And Bentgrass Lawns.
- Covers 5,000 Sq. Ft.
- Contains TRIMEC® Herbicide.

**ACTIVE INGREDIENTS:**

Dimethylamine salt of (+)-(R)-2-(2-methyl-4-chlorophenoxy)propionic acid.....	0.32%
Dimethylamine salt of 2,4-dichlorophenoxyacetic acid.....	0.18%
Dimethylamine salt of dicamba: 3,6-dichloro-o-anisic acid.....	0.08%

**INERT INGREDIENTS:**..... 99.42%

**TOTAL:**..... 100.00%

**THIS PRODUCT CONTAINS:**

5.40 lbs. or 0.27% (+)-(R)-2-(2-methyl-4-chlorophenoxy) propionic acid equivalent per ton.  
 3.00 lbs. or 0.15% 2,4-dichlorophenoxyacetic acid equivalent per ton.  
 1.40 lbs. or 0.07% 3,6-dichloro-o-anisic acid equivalent per ton.  
 Isomer Specific by AOAC Methods.  
 Contains the single isomer form of Mecoprop-p.  
 TRIMEC® is a registered trademark of PBI/Gordon Corporation.

**KEEP OUT OF REACH OF CHILDREN**

**CAUTION**

See Back Panel for Additional Precautionary Statements, First Aid and Directions for Use.



**NET WEIGHT 18 LBS. (8.16 KG)**

**33408**

**WEED & FEED**

**BROADLEAF WEEDS**

- Annual Yellow Sweetclover
- Aster
- Austrian Fieldgrass
- Bedstraw
- Beggarick
- Betony, Florida
- Bindweed, Field
- Bird Vetch
- Bitterweed
- Bittercress, Hairy
- Brier Wintercress
- Brickwallflower, Milkweed
- Broadleaf Plantain
- Broadleaf Dock
- Bristly Ox-tongue
- Broomsrape
- Buckhorn
- Buckhorn Plantain
- Bull Thistle
- Bull Thistle
- Bull Thistle
- Burncup
- Burned
- Burned
- Buttercup
- Buttercup
- Canada Thistle
- Carpetweed
- Catchweed Bedstraw
- Catsnip
- Catsnip
- Chickweed, Common
- Chickweed, Mouseear
- Chicory
- Chicory
- Clover, Common
- Clover, Hop
- Clover, Red
- Clover, Strawberry
- Clover, Sweet
- Clover, White
- Cockle
- Cocklebur
- Common Milkweed
- Creeping Jenny
- Cudweed
- Curl Dock
- Daisy, English
- Daisy, Fleabane
- Daisy, Oxeye
- Dandelion
- Dichondra
- Dogbane
- Dogbane
- Dollarweed
- False Dandelion
- False Sunflower
- Fiddleneck
- Florida Pusley
- Frenchweed
- Galinsoga
- Goldenrod
- Ground Ivy
- Gumweed
- Hairy Fleabane
- Hawweed
- Heartleaf
- Heartleaf Dymal
- Hedge Bindweed
- Hedge Mustard
- Henbit
- Hoary Cress
- Hoary Plantain
- Hoary Vervain
- Horsetail
- Jimsonweed
- Knapweed
- Kochia
- Lambsquarters
- Lespedeza
- Mallow
- Matchweed
- Mexicanweed
- Milk Vetch
- Morningglory
- Mouseear Hawkweed
- Mugwort
- Musk Thistle

**STOP! READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.**

**HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

**CAUTION:** Harmful if swallowed. Do not inhale dust. Do not allow contact with skin, eyes or clothing. When using this product, wear long-sleeved shirt, long pants, socks and shoes. After using this product, remove clothing and launder separately before reuse, and promptly and thoroughly wash hands and exposed skin with soap and water.

**FIRST AID**

- If In Eyes:**
  - Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing.
- If On Skin:**
  - Take off contaminated clothing.
  - Rinse skin immediately with plenty of water for 15-20 minutes.
- If Swallowed:**
  - Call a Poison Control Center or doctor immediately for treatment advice.
  - Have person sip a glass of water if able to swallow.
  - Do not induce vomiting unless told to do so by a Poison Control Center or doctor.

Have the product container or label with you when calling a Poison Control Center or doctor or going for treatment.

**ENVIRONMENTAL HAZARDS**

This product is toxic to aquatic invertebrates. Drift or runoff may adversely affect non-target plants. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area.

**DIRECTIONS FOR USE**

Sweeping any product that lands on a driveway, sidewalk, or street, back onto the treated area of the lawn or garden will help to prevent run off to water bodies or drainage.

**STORAGE AND DISPOSAL**

Do not contaminate water, food, or feed by storage and disposal. Do not allow people (other than applicator) or pets on treatment area during application. Do not enter treatment area until spray has dried or dust has settled.

**USE PRECAUTIONS**

Do not use on Dichondra. Do not overapply. Under conditions of adequate moisture, high temperature and high humidity, over-application may cause minor transitory discoloration on St. Augustinegrass.

Do not use this product where desirable Clovers are present or allow the material to wind drift onto flowers, vegetables or ornamental shrubbery. Do not apply in areas underlaid by roots of desirable trees or shrubs. Apply when air is calm. Do not apply to newly seeded lawns. During application, close spreader openings when turning.

**WHEN TO APPLY:**

**H-YIELD® WEED & FEED** is a Trimec complex, a combination of three proven weed killers - 2,4-D, Mecoprop and Dicamba. Together, they control a wide range of Lawn Weeds, including Dandelion, Chickweed, Knotweed, Plantain, Henbit, Spurge and many others commonly found in Home Lawns. In addition, a 15-0-10 fertilizer provides a supply of nutrients. Possible re-treatment may be necessary if (a) Lawns is heavily overrun by Weeds, or, (b) if adverse low moisture conditions prevail and Weeds are in a state of poor growth. A second application may be made in approximately 30 days.

The maximum application rate to turf is 0.25 pounds of 2,4-D acid equivalent per acre per application per site. The maximum number of broadcast applications per treatment site is 2 per year.

1. Mow Lawn to normal height 1 to 2 days before application.

2. Water Lawn thoroughly at least 1 day before application to sustain moisture until the next watering (See Step 4).

3. Apply when Weeds are young and actively growing, preferably in the morning when dew is on the Grass. If Grass is not moist at time of application, sprinkle lightly with water to hold the material and prevent dust drift onto non-target plants.

4. Do not wash from Weed leaves for 1 to 2 days after application. At this time, a thorough watering should be made.



This application map estimates the schedule of fertilizer applications and should be used as a guideline only. Abnormal climatic conditions can affect your schedule.		
April through Mid-June	Mid-March through May	January through April
Northern Zone	Transition Zone	Southern Zone

SPREADER SETTINGS	
MODEL	3.2
fertiHome® By SPYKER	17
fertiHome® PLUS IRON BROADCAST	17
fertiHome® PLUS IRON PROF. BROADCAST	4
REPUBLIC BROADCAST	5
REPUBLIC DROP	11
EARTHWAY DROP	11

When used as directed this product will provide 0.5 lb of nitrogen per 1,000 square feet. Another Hi-Yield fertilizer product may be used 30 days after application to supplement this application.

**15-0-10 GUARANTEED ANALYSIS**

Total Nitrogen (N).....15%  
 2.0% Urea Nitrogen  
 4.5% Water Insoluble Nitrogen  
 8.5% Ammoniacal Nitrogen

Dewfall From: Ammonium Sulfate, Urea, Urea Formaldehyde, Mixture of Potash Soluble Potash (K<sub>2</sub>O).....10%

\* 4.5% Slowly Available Nitrogen from Urea Formaldehyde

**LIMITED WARRANTY AND DISCLAIMER**

The Manufacturer warrants only that the chemical composition of this product conforms to the ingredient statement given on the label, and that the product is reasonably suited for the labeled use when applied according to the Directions For Use.

THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY OTHER EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE EXPRESSLY DISCLAIMED. This Limited Warranty does not extend to the use of the product inconsistent with label instructions, warnings or cautions, or to use of the product under abnormal conditions such as drought, excessive rainfall, tomatoes, hurricanes, etc. These factors are beyond the control of the Manufacturer or the Seller. Any damages arising from a breach of the Manufacturer's Warranty shall be limited to direct damages, and shall not include indirect or consequential damages such as loss of profits or values, except as otherwise provided by law.

The terms of this Limited Warranty and Disclaimer cannot be varied by any written or verbal statements or agreements. No employee or agent of the Seller is authorized to vary or exceed the terms of this Limited Warranty and Disclaimer in any manner.

Produced For:  
**VOLUNTARY PURCHASING GROUPS, INC.**  
 230 FM 87  
 BONHAM, TEXAS 75418  
 Visit Us At: www.vh-yield.com  
 EPA Reg. No. 227-661-7401 • EPA Est. No. 7401-TX-1



50M-1-19-2

**BROADLEAF WEEDS**

- Mustard
- Narrowleaf Plantain
- Narrowleaf Vetch
- Nettle
- Orange Hawkweed
- Oriental Cocklebur
- Oxalis
- Parsley-Fiert
- Parsnip
- Peawort
- Pennygrass
- Pennywort
- Pepperglass
- Pepperweed
- Pigweed
- Pinewoods Bedstraw
- Plains Coreopsis (Tickseed)
- Plantain
- Poison Ivy
- Poison Oak
- Poisoned
- Portulac
- Prairie Sunflower
- Prickly Lettuce
- Prickly Sida
- Prostrate Knotweed
- Prostrate Plantain
- Prostrate Spurge
- Prostrate Vervain
- Puncturevine
- Purslane, Common
- Ragweed
- Red Sorrel
- Redroot Pigweed
- Redstem Filaree
- Rough Chickweed
- Rough Fleabane
- Russian Figweed
- Russian Thistle
- Scarlet Pimpernel
- Scotch Thistle
- Sheep Sorrel
- Shepherdspurse
- Slender Plantain
- Smallflower Galinsoga
- Smooth Dock
- Smooth Figweed
- Sorrel
- Sowthistle
- Spatismeedles
- Speedwell
- Spotted Spurge
- Spiny Amaranth
- Spiny Cocklebur
- Spiny Sowthistle
- Spotted Catsear
- Spurred
- Singing Nettle
- Strawberry, India Mock
- Tail Nettle
- Tall Vervain
- Tansy Ragwort
- Tansy Mustard
- Thistle
- Trailing Crownvetch
- Tumble Mustard
- Tumble Pigweed
- Valeriate
- Venice Mallow
- Virginia Buttonweed
- Virginia Creeper
- Virginia Peppweed
- Water Pennywort
- Wayleaf Bullthistle
- Western Clematis
- Western Salsify
- White Mustard
- Wild Aster
- Wild Buckwheat
- Wild Carrot
- Wild Four-O-Clock
- Wild Garlic
- Wild Geranium
- Wild Lettuce
- Wild Margold
- Wild Mustard
- Wild Onion
- Wild Parsnip
- Wild Radish
- Wild Rape
- Wild Strawberry
- Wild Sweet Potato
- Wild Vetch
- Woodsorel
- Woolly Croton
- Woolly Morningglory
- Woolly Plantain
- Wormseed
- Yarrow
- Yellow Rocket
- Yellowflower Pepperweed
- and other Broadleaf Weeds